



XIIIth Meeting on Lattice Parton Physics
from Large Momentum Effective Theory
(LaMET 2026)



Probing Instanton Dynamics in the Pion Vector Form Factor with Gradient Flow


— Vaibhav Chahar, Piotr Korcyl —

Jagiellonian University, Poland

LaMET 2026

Introduction

- QCD is a non-Abelian gauge theory based on the SU(3) color symmetry group. It describes interactions between quarks and gluons.
- The QCD vacuum is not empty, as it contains topologically non-trivial gluon field configurations.
- Instantons are localized, classical solutions of the Euclidean Yang–Mills equations.
- They describe **tunneling** between distinct classical QCD vacua.



R. Jackiw and C. Rebbi, *Phys. Rev. Lett.* 37, 172 (1976)

Wilson Flow (Gradient Flow)

- The QCD vacuum on the lattice is full of UV noise, tiny fluctuations that obscure physical, topological structures.
- Wilson flow is a process that evolves lattice gauge fields $U_\mu(\mathbf{x})$ along a continuous “flow time” t by solving:

$$\frac{dU_\mu(\mathbf{x},t)}{dt} = -g_0^2 \left\{ \frac{\partial S_g[U(t)]}{\partial U_\mu(\mathbf{x},t)} \right\} U_\mu(\mathbf{x},t), \quad U_\mu(\mathbf{x},0) = U_\mu(\mathbf{x}),$$

- The gauge fields become smoother as t increases, with small-scale noise diffused away.
- Applications in Lattice Field Theory:
 - Topological Charge and Susceptibility
 - Scale Setting
 - Improve Signal to noise

Wilson Flow and Instantons


- Each instanton carries a topological charge corresponding to tunneling between degenerate QCD vacua.

$$Q = \frac{1}{32\pi^2} \int d^4x G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a$$

- At **small flow times**, UV fluctuations are suppressed, but physical long-distance structures remain intact.
- At **moderate flow**, instanton-like objects appear as **coherent lumps** in the topological charge.
- If the flow time is too long, instantons may shrink and annihilate (over-smoothing), so an **optimal range of flow time** is chosen.
- In this work, we study hadronic matrix elements directly at finite Wilson flow time $t > 0$, and understand how they change as the flow time is varied.

Instantons and Pion Form Factors

- The **Instanton Liquid Model (ILM)** provides a mathematical framework to describe this complex vacuum by modeling it as an ensemble of interacting instantons and anti-instantons.
- One particularly insightful observable is the **pion electromagnetic form factor**, which probes the internal structure and dynamics of the pion as a bound state of quarks.
- The Pure Instanton Liquid Model gives a higher value of the pion form factor and does not agree with lattice qcd and experiments.
- We study the pion form factor for different values for t .



Pion Form factors from Lattice

$$O_\pi(x,t) = \bar{d}(x,t) \gamma_5 u(x,t),$$

$$O_\pi^\dagger(x,t) = \bar{u}(x,t) \gamma_5 d(x,t).$$


$$C_\pi^{2\text{pt}}(t, \mathbf{p}) = \sum_{\mathbf{x}} e^{-i\mathbf{p}\cdot\mathbf{x}} \langle 0 | O_\pi(\mathbf{x}, t) O_\pi^\dagger(0, 0) | 0 \rangle.$$

$$C_\pi^{3\text{pt}}(t, t_{\text{sep}}, \mathbf{p}_f, \mathbf{p}_i) = \sum_{\mathbf{x}, \mathbf{y}} e^{-i\mathbf{p}_f\cdot(\mathbf{x}-\mathbf{y})} e^{-i\mathbf{p}_i\cdot\mathbf{y}} \langle 0 | O_\pi(\mathbf{x}, t_{\text{sep}}) J_\mu(\mathbf{y}, t) O_\pi^\dagger(0, 0) | 0 \rangle.$$

$$R(t_{\text{sink}}, \tau, \vec{p}, \vec{p}') = \frac{C_{3\text{pt}}^O(\tau, \vec{p}', \vec{p})}{C_{2\text{pt}}(t_{\text{sink}}, \vec{p}')} \sqrt{\frac{C_{2\text{pt}}(t_{\text{sink}} - \tau, \vec{p}) C_{2\text{pt}}(\tau, \vec{p}') C_{2\text{pt}}(t_{\text{sink}}, \vec{p})}{C_{2\text{pt}}(t_{\text{sink}} - \tau, \vec{p}) C_{2\text{pt}}(\tau, \vec{p}) C_{2\text{pt}}(t_{\text{sink}}, \vec{p})}}.$$

$$R(t_{\text{sink}}, \tau, p', p) = \frac{\langle \pi(p') | O(\tau) | \pi(p) \rangle}{2}$$

$$2\sqrt{E_{\vec{p}'} E_{\vec{p}}} \times Z_V \times \langle \pi(\vec{p}') | \mathcal{O}_V^\mu(0) | \pi(\vec{p}) \rangle = (p_f + p_i)_\mu \times f_{\pi\pi}(-q^2)$$




Lattice Setup

- The study use publicly available PACS-CS gauge fields ensemble.
- Fermionic part o(a)-improved wilson action with $N_f = 2 + 1$ dynamical quarks and gauge part is Iwasaki gauge action.
- Bare parameters for the ensemble



β	a [fm]	k_1	L/a	T/a	c_{sw}	N_G	M_π	Z_v
1.96	0.0907(13)	0.13754	32	64	1.715	200	409.7(7)	0.7354(37)

Procedure




Setting up the bare parameters

- Preliminary: set $c_{\text{sw}} = 1.0$. A more precise analysis requires tuning c_{sw} and c_A for example using PCAC relations.
- Rather than performing ADDITIONAL simulations at vanishing quark mass, we attempt to introduce a massive renormalization and improvement scheme, extracting the relevant parameters directly from our available ensemble at $M_\pi \simeq 400$ MeV.



Setting up the bare parameters

We see a significant difference between the form factors at tree-level and flow time = $5t_0$



Setting up the bare parameters

- PCAC relation

$$\langle \partial_\mu A_\mu(x) \mathcal{O}(y) \rangle = 2m \langle P(x) \mathcal{O}(y) \rangle \quad \text{where, } A_\mu(x) = \bar{u}(x) \gamma_\mu \gamma_5 d(x), P(x) = \bar{u}(x) \gamma_5 d(x)$$

For simplicity, $\mathcal{O}(y) = P^\dagger(y)$

$$\frac{1}{2} (\partial_\mu^* + \partial_\mu) \langle A_\mu^I(x) P^\dagger(y) \rangle = 2m \langle P(x) P^\dagger(y) \rangle$$

$$A_\mu^I(x) = A_\mu(x) + c_A \frac{1}{2} (\partial_\mu^* + \partial_\mu) P(x)$$

We set $\mu = 0$

$$\vec{\partial}_0 \langle \{ A_0(x) + c_A \vec{\partial}_0 \} P(x) \rangle P^\dagger(y) = 2m \langle P(x) P^\dagger(y) \rangle$$

- With symmetric derivative $\overleftrightarrow{\partial}_0 = \frac{1}{2} (\partial_0^* + \partial_0)$

$$\vec{\partial}_0 \langle A_0(x) P^\dagger(y) \rangle + c_A \overleftrightarrow{\partial}_0 \vec{\partial}_0 \langle P(x) P^\dagger(y) \rangle = 2m \langle P(x) P^\dagger(y) \rangle$$

- Now we have ratio


$$m(x, y; c_A, c_{SW}, m_\pi) = \frac{\overleftrightarrow{\partial}_0 A_d \langle A_0(x) P^\dagger(y) \rangle + c_A \overleftrightarrow{\partial}_0 \vec{\partial}_0 \langle P(x) P^\dagger(y) \rangle}{\langle P(x) P^\dagger(y) \rangle}$$

Setting up the bare parameters


$$m(x, y; c_A, c_{SW}, m_\pi) = \frac{\overline{\partial}_0 \langle A_0(x) P^\dagger(y) \rangle + c_A \overline{\partial}_0 \overline{\partial}_0 \langle P(x) P^\dagger(y) \rangle}{\langle P(x) P^\dagger(y) \rangle}.$$

- The ratio is plotted with c_A
- Choose the points in the cone which lies on the hyper diagonal of the lattice.
- The cone is defined using **physical distances**.
- Then choose the points which lies inside the shaded frustum.
- Larger the angle θ , the more points to consider
- Calculate the variance of $m(y; c_{SW}, c_A)$ using points lies in the frustum various c_{sw} and M_π .


$$\text{Var}(m) = \frac{1}{N-1} \sum_{y \in \mathcal{C}} (m(y) - \bar{m})^2$$



ca vs var for csw 1.720





ca vs var for csw 1.730



Setting up the bare parameters



- Once we fix c_{sw} we will know all necessary parameters and could estimate the form factor.
- The algorithm can be applied at $t \triangleright 0$.



Preliminary results for parameter tuning at $t = 0$, for 20 gauge configurations.

Setting up the bare parameters

- Can matrix elements at large flow time be extracted using the same procedure as at $t = 0$?
- Model calculations suggest that the standard extraction procedure remains valid at finite flow time.
- After tuning κ for each flow time, the pion mass is consistent across flows. However, excited-state contributions are enhanced at larger flow times.



Summary and Outlook

- Tune the parameters c_{sw} , κ
- In future, form factor will be calculated with twisted boundary conditions to get access to more momenta between the integer values.
- After tuning the parameters, calculation of form factors with higher values of flow time will be used.
- Extrapolation to $a \rightarrow 0$
- Use more lattice ensembles for broader result.

References

1. Chu, M.-C., Grandy, J. M., Huang, S., & Negele, J. W. **Evidence for the Role of Instantons in Hadron Structure from Lattice QCD**. *Physical Review D*, **49**, 6039–6050 (1994). arXiv:hep-lat/9312071.
2. arXiv: **hep-lat/9208030**. Preprint available at: <https://arxiv.org/abs/hep-lat/9208030>
3. arXiv: **hep-lat/9211019**. Preprint available at: <https://arxiv.org/abs/hep-lat/9211019>
4. arXiv: **hep-lat/9312071**. Preprint available at: <https://arxiv.org/abs/hep-lat/9312071>

Thank You